

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number
WO 01/96314 A1

(51) International Patent Classification⁷: **C07D 239/48**,
401/04, A01N 43/54

(21) International Application Number: **PCT/EP01/06565**

(22) International Filing Date: **11 June 2001 (11.06.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/211,262 13 June 2000 (13.06.2000) US
60/231,632 11 September 2000 (11.09.2000) US

(71) Applicant: **BASF AKTIENGESELLSCHAFT**
[DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors: **PEES, Klaus-Juergen**; Soonwaldstrasse 9,
55192 Mainz (DE). **PFRENGLE, Waldemar**; Junker-
muehle 1, 55444 Siebersbach (DE). **HEFFERNAN,**
Gavin; 616 East Sixth Street, Florence, NJ 08518 (US).

(74) Common Representative: **BASF AKTIENGE-**
SELLSCHAFT; 67056 Ludwigshafen (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **FUNGICIDAL 5-PHENYL SUBSTITUTED 2-(CYANOAMINO) PYRIMIDINES**



(I)


(57) Abstract: Pyrimidines of formula (I) in which R¹ represents hydrogen or alkyl, haloalkyl, alkenyl, alkynyl, alkadienyl, alkoxy, cycloalkyl, phenyl, or 5- or 6-membered heteroaryl or 5- or 6-membered heterocyclyl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or tri-alkyl-silyl, formyl or alkoxycarbonyl, wherein R¹ groups are unsubstituted or substituted as defined in the specification; R² represents phenyl, cycloalkyl or 5- or 6-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, which are unsubstituted or substituted; R³ represents hydrogen, halogen or alkyl, alkoxy, alkylthio, alkylamino or dialkylamino; which are unsubstituted or substituted; R⁴ represents hydrogen or alkyl, alkenyl or alkynyl; which are unsubstituted or substituted; and X represents O, S, NR⁵ or a single bond, wherein R⁵ represents hydrogen or alkyl; or R¹ and R⁵

together with the interjacent nitrogen atom form a heterocyclic ring; processes and intermediates for preparing these compounds, to compositions comprising them and to their use for controlling harmful fungi.

WO 01/96314 A1

FUNGICIDAL 5-PHENYL SUBSTITUTED 2-(CYANOAMINO) PYRIMIDINES

The present invention relates to pyrimidines of formula I



10

in which

R¹ represents hydrogen or

15

C₁-C₁₀-alkyl, C₁-C₁₀-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₄-C₈-alkadienyl, C₁-C₁₀-alkoxy, C₃-C₈-cycloalkyl, phenyl, or

5- or 6-membered heteroaryl or 5- or 6-membered heterocyclyl, containing one to four nitrogen atoms or one to three

20

nitrogen atoms and one sulfur or oxygen atom, or

tri-C₁-C₆-alkyl-silyl, formyl or C₁-C₁₀-alkoxycarbonyl;

wherein R¹ groups are unsubstituted or substituted by one to

25

three groups R^a

R^a halogen, nitro, cyano, hydroxy or

30

C₁-C₁₀-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₁-C₁₀-haloalkyl, C₃-C₆-halocycloalkyl, C₁-C₁₀-alkoxy, C₁-C₁₀-haloalkoxy, C₁-C₁₀-haloalkoxy, C₁-C₆-alkoxycarbonyl, tri-C₁-C₄-alkylsilyl, phenyl, halo- or dihalo-phenyl or 5- or 6-membered heteroaryl, containing one to four

35

nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom;

R² represents phenyl, C₃-C₆-cycloalkyl or 5- or 6-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, which are

40

unsubstituted or substituted by one to three groups R^a;

R³ represents hydrogen, halogen or

45

C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy, C₁-C₁₀-alkylthio, C₁-C₁₀-alkylamino or di-C₁-C₁₀-alkylamino; which are unsubstituted or substituted by one to three groups R^a;

R⁴ represents hydrogen or

C₁-C₁₀-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl; which are unsubstituted or substituted by one to three groups R^a; and

5

X represents O, S, NR⁵ or a single bond, wherein R⁵ represents hydrogen or C₁-C₁₀-alkyl; or

10 R¹ and R⁵ together with the interjacent nitrogen atom form a heterocyclic ring.

Moreover, the invention relates to processes and intermediates for preparing these compounds, to compositions comprising them and to their use for controlling harmful fungi.

15

In Vestn. Slov. Kem. Drus. (1986), 33(3), 353-66 (ISSN: 0560-3110, CAN 107:39701) it is disclosed that the reaction of N-pyrimid-2-ylformamide oximes with N,N-dimethylformamide diethyl acetal yields 2-(N-cyano-N-ethylamino)pyrimidines. In J. Org. Chem. 39 (9) 1256-1252 (1974) N-glycosylated 2-(N-cyanoamino) pyrimidines are disclosed and in US 4,711,959 a process for the preparation of 2-(N-cyanoamino) pyrimidines is described.


25 It is an object of the present invention to provide fungicidal compounds having improved activity.

We have found that this object is achieved by the compounds defined at the outset. Furthermore, we have found processes for their preparation, compositions comprising them and methods for controlling harmful fungi using the compounds I.

30

Compounds of formula I wherein R⁴ is an optionally substituted alkyl, alkenyl or alkynyl group as defined above may be obtained by treating a compound of the formula II

35



II

40

in which R¹ through R³ and X are as defined in formula I; with a strong base and an alkylation agent of formula III

45

R⁴-Y

III

3

in which R^4 is C_1 - C_6 -alkyl, C_1 - C_6 -alkenyl or C_1 - C_6 -alkynyl; which are unsubstituted or substituted by one to three groups R^a , and Y represents a nucleophilic replaceable leaving group, preferably a halogen atom, in particular a iodine atom.

5

Compounds of formula II are known for example from US 5,593,996, WO-A 98/46608, FR-A 2,765,875, WO-A 99/41255 or WO-A 99/48893.


The reaction between the triazolopyrimidines of formula II, the strong base and the alkylation agent of formula III is preferably carried out in the presence of an inert solvent. Suitable solvents include ethers, such as dioxane, diethyl ether and tetrahydrofuran, halogenated hydrocarbons such as dichloromethane, amides, such as dimethylformamide or N-methylpyrrolidone and aromatic hydrocarbons, for example toluene or mixtures of these solvents. The reaction is suitably carried out at a temperature in the range from -78°C to 100°C , the preferred reaction temperature is from 10°C to 80°C , particular at ambient temperature.

20

Suitable strong bases include metal hydrides, such as sodium hydride, potassium hydride or calcium hydride, and metal amides, such as sodium amide, potassium amide, lithium diisopropylamide or potassium hexamethyldisilazide, and metal alkanes such as methyllithium, n-butyllithium or tert-butyllithium.

30

Furthermore, the compounds of formula I wherein R^4 is an optionally substituted alkyl, alkenyl or alkynyl group may be prepared by reacting a N-pyrimid-2-ylformamide oxime of formula IV



in which R^1 through R^3 and X are as defined in formula I; with a N,N-dimethylformamide dialkyl acetate of formula V



40


in which R^4 is C_1 - C_6 -alkyl, C_1 - C_6 -alkenyl or C_1 - C_6 -alkynyl; which are unsubstituted or substituted by one to three groups R^a .

The reaction between the compounds of formula IV and the compounds of formula V can be carried out analogously to the reaction described in Vestn. Slov. Kem. Drus. (1986), 33(3), 353-66.

4

Compounds of formula I wherein R^4 is hydrogen can preferably be prepared by treating sulfones of formula VI

5



VI

in which R^1 through R^3 and X are as defined in formula I and R^6 is C_1 - C_6 -alkyl or C_1 - C_6 -haloalkyl; with cyanamide or with a metal salt of cyanamide. The use of a base and/or a solvent can be advantageous.

This process is preferably carried out in the presence of an inert solvent. Suitable solvents include aromatic hydrocarbons, such as, for example toluene or xylene, chlorinated hydrocarbons, such as, for example methylene chloride, chloroform, a chlorobenzene, ketones, such as, for example acetone, methyl ethyl ketone, methyl isopropyl ketone or methyl isobutyl ketone, nitriles, such as, for example acetonitrile or propionitrile ethers, such as, for example diethyl ether, diisopropyl ether, methyl tert-butylether, dimethoxyethane, tetrahydrofuran or dioxane, amides, such as, for example, dimethylacetamide or diethylacetamide, sulfoxides, such as, for example dimethylsulfoxide or sulfolane, or mixtures thereof.

The use of a base can be advantageous in this reaction. Suitable bases include alkali metal hydrides and earth alkaline metal hydrides, such as, for example, sodium, potassium or calcium hydrides, alkali metal hydroxides and alkaline earth metal hydroxides, such as, for example, sodium, potassium or calcium hydroxides, alkali metal carbonates and alkaline earth metal carbonates, such as, for example sodium carbonate, potassium carbonate or calcium carbonate, alkali metal bicarbonates and alkaline earth metal bicarbonates, such as sodium bicarbonate, potassium bicarbonate or calcium bicarbonate, metal amides, such as, for example sodium amide, potassium amide, lithium diisopropylamide or potassium hexamethyldisilazide, metal alkanes, such as for example methyl lithium, n-butyl lithium or tert-butyl lithium or aprotic amines, such as, for example pyridine, tributylamine, N,N-dimethylbenzylamine or diazobicycloundecene.

Various qualities of cyanamide may be employed for the process. The use of an aqueous solution of cyanamide may be preferred for practical reasons. The use of metal salts of cyanamide, potassium cyanamide, dipotassium cyanamide or calcium cyanamide is also possible.

Dependant on the used cyanamide or salt of, cyanamide and dependant on the base an appropriate solvent is employed.

The reaction is suitable carried out at a temperature in the range from -78°C to reflux temperature, the preferred reaction temperature is from 0°C to 150°C, particular at ambient temperature.

In general 1 to 3 equivalents, preferably 1.5 to 2.5 equivalents of base are employed per equivalent of sulfone of the formula VI.

Generally 2 to 6 equivalents, preferably 3 to 5 equivalents of cyanamide or salt of cyanamide are employed per equivalent of the sulfone of the general formula VI.

15

Compounds of formula I wherein R⁴ is C₁-C₆-alkyl, C₁-C₆-alkenyl or C₁-C₆-alkynyl which are unsubstituted or substituted by one to three groups R^a may be prepared by alkylation of compounds of formula I wherein R⁴ is hydrogen with an alkylation agent of formula

20 III.

The use of a base can be advantageous in this reaction. Suitable bases include alkali metal hydrides and earth alkaline metal hydrides, such as, for example, sodium, potassium or calcium hydrides, alkali metal hydroxides and alkaline earth metal hydroxides, such as, for example, sodium, potassium or calcium hydroxides, alkali metal carbonates and alkaline earth metal carbonates, such as, for example sodium carbonate, potassium carbonate or calcium carbonate, alkali metal bicarbonates and alkaline earth metal bicarbonates, such as sodium bicarbonate, potassium bicarbonate or calcium bicarbonate, metal amides, such as, for example sodium amide, potassium amide, lithium diisopropylamide or potassium hexamethyldisilazide, metal alkanes, such as for example methyl lithium, n-butyl lithium or tert-butyl lithium or aprotic amines, such as, for example pyridine, tributylamine, N,N-dimethylbenzylamine or diazobicycloundecene.

The alkylation is preferably carried out in the presence of an inert solvent. Suitable solvents include aromatic hydrocarbons, such as, for example toluene or xylene, chlorinated hydrocarbons, such as, for example methylene chloride, chloroform, a chlorobenzene, ketones, such as, for example acetone, methyl ethyl ketone, methyl isopropyl ketone or methyl isobutyl ketone, nitriles, such as, for example acetonitrile or propionitrile ethers, such as, for example diethyl ether, diisopropyl ether, methyl tert-butylether, dimethoxyethane, tetrahydrofuran or dioxane, amides, such as, for example, dimethylacetamide or diethylacetamide, sulfoxi-

6

des, such as, for example dimethylsulfoxide or sulfolane, or mixtures thereof.


The reaction is suitably carried out at a temperature in the range from -78°C to reflux temperature, the preferred reaction temperature is 0°C to 150°C , particular ambient temperature.

In general 0.8 to 5 equivalents, preferably 0.8 to 4.5 equivalents of the alkylation agent of the formula III are employed per equivalent of the compound of formula I.

Usually 0.8 to 3 equivalents, preferably 0.8 to 2.5 equivalents of base are employed per equivalent of the compound of formula I.

It is also possible to prepare compounds of formula I wherein R^4 is $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-alkenyl}$ or $\text{C}_1\text{-C}_6\text{-alkynyl}$ which are unsubstituted or substituted by one to three groups R^a by reacting a sulfone of formula VI


20



VI

in which R^1 through R^3 and X are as defined in formula I and R^6 is $\text{C}_1\text{-C}_6\text{-alkyl}$ or $\text{C}_1\text{-C}_6\text{-haloalkyl}$; with an alkylated cyanamide of formula VII

30



VII

wherein R^4 is $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-alkenyl}$ or $\text{C}_1\text{-C}_6\text{-alkynyl}$ which are unsubstituted or substituted by one to three groups R^a . The use of a base and/or a solvent can be advantageous.

35

Suitable bases and solvents are such as listed at the reaction with cyanamide.

The reaction is suitable carried out at a temperature in the range of from -78°C to reflux temperature, the preferred reaction temperature is from 0°C to 150°C , particular at ambient temperature.

The reaction is in general carried out under usual pressure.

45


In general 1 to 3 equivalents preferably 1.5 to 2.5 equivalents of base are employed per equivalent of the sulfone of formula VI.

7

Usually 2 to 6 equivalents, preferably 3 to 5 equivalents of alkylated cyanamide of formula VII are employed per equivalents of the sulfone of formula VI.

- 5 Sulfones of the formula VI are obtained by reacting 2-thiopyrimidinederivatives of the formula VIII

10




VIII

in which the variables are as defined in formula VI; with oxidizing agents, such as, for example m-chloroperbenzoic acid, per
15 acetic acid, trifluoro per acetic acid, chlorine water, hypocho-
rous acid or metal salt solutions thereof in water or hydrogen
peroxide, if appropriate in presence of a catalyst, such as for
example wolframate.

- 20 If appropriate solvents, such as for example, methylene chloride, chloroform, carbontetrachloride, 1,2-dichloroethane or chloroben-
zene are used at temperatures of -20°C to reflux.

The 2-thiopyrimidine derivatives of the formula VIII may be ob-
25 tained when 6-halo-2-thiopyrimidines of formula IX


30



IX

in which the substituents are as before mentioned and "Hal" deno-
tes halogen; are reacted with a nucleophile of formula X

35




X

wherein R¹ and X are as defined in formula I, if appropriate in
the presence of a suitable base and if appropriate in an organic
40 solvent. The solvents and bases employed are similar to those
mentioned for the preparation of the compounds of formula I.

6-halo-2-thiopyrimidines of formula IX are known in the art or
may be prepared according following reaction sequence:

45



(R², R³ and R⁶ are as defined above and R is an alkyl group) The reaction conditions are in general known in the art.

Base catalyzed reaction of dialkylmalonate with thiourea affords 2-thiobarbituric acid XII which may be selectively alkylated on sulfur to yield XIII.

Halogenation, preferably chlorination or bromination, especially chlorination, with for example phosphorous oxychloride or phosphorous oxybromide in the presence of a tertiary amine base then affords the dihalo derivative XIV.

Subsequent introduction of the radical R³, if appropriate, via nucleophilic substitution affords the 6-halo-2-thiopyrimidine of formula IX.

Sulfones of formula VI and 2-thiopyrimidine derivatives of formula VIII are novel.

In the symbol definitions given in the formulae above, collective terms were used which generally represent the following substituents:

- halogen: fluorine, chlorine, bromine and iodine;

- C₁-C₆-alkyl and the alkyl moieties of C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylamino, di-C₁-C₆-alkylamine or

C₁-C₆-alkoxycarbonyl: saturated, straight-chain or branched hydrocarbon radicals having 1 to 6 carbon atoms, preferably C₁-C₄-alkyl, such as methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl; or pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-di-methylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl; preferably ethyl or methyl;

- C₁-C₆-haloalkyl and the haloalkyl moieties of C₁-C₆-haloalkoxy:
15 straight-chain or branched alkyl groups having 1 to 6 carbon atoms (as mentioned above), where the hydrogen atoms in these groups may be partially or fully replaced by halogen atoms as mentioned above, for example C₁-C₂-haloalkyl, such as chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, 20 fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 25 2,2,2-trichloroethyl and pentafluoroethyl; preferably 2,2,2-trifluoroethyl or 1,1,1-trifluoroprop-2-yl;

- C₃-C₈-cycloalkyl: monocyclic, saturated hydrocarbon radicals having 3 to 6 or 8 carbon ring members, for example cyclopropyl, 30 cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, preferably 5 to 7 carbon atoms, in particular cyclopentyl being optionally substituted by one or more halogen atoms, nitro, cyano, C₁-C₆-alkyl or C₁-C₆-alkoxy.

35 - C₂-C₄-alkenyl: unsaturated, straight-chain or branched hydrocarbon radicals having 2 to 4 carbon atoms and a double bond in any position, for example ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl and 40 2-methyl-2-propenyl; preferably allyl or 2-methylallyl.

- C₂-C₄-haloalkenyl and the haloalkenyl moieties of C₂-C₄-haloalkenyloxy: unsaturated, straight-chain or branched hydrocarbon radicals having 2 to 4 carbon atoms and a double bond 45 in any position (as mentioned above), where the hydrogen atoms in these groups may be partially or fully replaced by halogen atoms

10

as mentioned above, in particular by fluorine, chlorine and bromine;

- C₂-C₄-alkynyl: straight-chain or branched hydrocarbon radicals
5 having 3 to 4 carbon atoms and a triple bond in any position, for example ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne and 1-methyl-2-propynyl;

- C₃-C₄-haloalkynyl and the haloalkynyl moieties of
10 C₂-C₄-haloalkynoxy: unsaturated, straight-chain or branched hydrocarbon radicals having 2 to 4 carbon atoms and a triple bond in any position (as mentioned above), where the hydrogen atoms in these groups may be partially or fully replaced by halogen atoms as mentioned above, in particular by fluorine, chlorine and
15 bromine;

5-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom: 5-membered heteroaryl groups which, in addition to carbon atoms, may
20 contain one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom as ring members, for example 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl,
25 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-imidazolyl, 4-imidazolyl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,4-triazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-thiadiazol-2-yl and 1,3,4-triazol-2-yl;

30

6-membered heteroaryl, containing one to four nitrogen atoms: 6-membered heteroaryl groups which, in addition to carbon atoms, may contain one to three or one to four nitrogen atoms as ring members, for example 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, 1,3,5-triazin-2-yl and 1,2,4-triazin-3-yl; preferred hetaryl moieties are pyridyl, pyrimidyl, pyrazolyl or thienyl.

40 5- or 6-membered heterocyclyl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, for example 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydrothienyl, 3-tetrahydrothienyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 3-isoxazolidinyl, 4-isoxazolidinyl, 5-isoxazolidinyl,
45 3-isothiazolidinyl, 4-isothiazolidinyl, 5-isothiazolidinyl, 3-pyrazolidinyl, 4-pyrazolidinyl, 5-pyrazolidinyl, 2-oxazolidinyl, 4-oxazolidinyl, 5-oxazolidinyl, 2-thiazolidinyl, 4-thiazolidinyl,

5-thiazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 1,2,4-oxadiazolidin-3-yl, 1,2,4-oxadiazolidin-5-yl, 1,2,4-thiadiazolidin-3-yl, 1,2,4-thiadiazolidin-5-yl, 1,2,4-triazolidin-3-yl, 1,3,4-oxadiazolidin-2-yl, 1,3,4-thiadiazolidin-2-yl, 1,3,4-triazolidin-2-yl, 2,3-dihydrofur-2-yl, 2,3-dihydrofur-3-yl, 2,4-dihydrofur-2-yl, 2,4-dihydrofur-3-yl, 2,3-dihydrothien-2-yl, 2,3-dihydrothien-3-yl, 2,4-dihydrothien-2-yl, 2,4-dihydrothien-3-yl, 2-pyrrolin-2-yl, 2-pyrrolin-3-yl, 3-pyrrolin-2-yl, 3-pyrrolin-3-yl, 2-isoxazolin-3-yl, 3-isoxazolin-3-yl, 4-isoxazolin-3-yl, 2-isoxazolin-4-yl, 3-isoxazolin-4-yl, 4-isoxazolin-4-yl, 2-isoxazolin-5-yl, 3-isoxazolin-5-yl, 4-isoxazolin-5-yl, 2-isothiazolin-3-yl, 3-isothiazolin-3-yl, 4-isothiazolin-3-yl, 2-isothiazolin-4-yl, 3-isothiazolin-4-yl, 4-isothiazolin-4-yl, 2-isothiazolin-5-yl, 3-isothiazolin-5-yl, 4-isothiazolin-5-yl, 2,3-dihydropyrazol-1-yl, 2,3-dihydropyrazol-2-yl, 2,3-dihydropyrazol-3-yl, 2,3-dihydropyrazol-4-yl, 2,3-dihydropyrazol-5-yl, 3,4-dihydropyrazol-1-yl, 3,4-dihydropyrazol-3-yl, 3,4-dihydropyrazol-4-yl, 3,4-dihydropyrazol-5-yl, 4,5-dihydropyrazol-1-yl, 4,5-dihydropyrazol-3-yl, 4,5-dihydropyrazol-4-yl, 4,5-dihydropyrazol-5-yl, 2,3-dihydrooxazol-2-yl, 2,3-dihydrooxazol-3-yl, 2,3-dihydrooxazol-4-yl, 2,3-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 3,4-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1,3-dioxan-5-yl, 2-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothienyl, 3-hexahydropyridazinyl, 4-hexahydropyridazinyl, 2-hexahydropyrimidinyl, 4-hexahydropyrimidinyl, 5-hexahydropyrimidinyl, 2-piperazinyl, 1,3,5-hexahydro-triazin-2-yl and 1,2,4-hexahydrotriazin-3-yl; preferred heterocyclyl groups are pyrrolidinyl, pyrazolidinyl, piperidinyl, piperazinyl or morpholin-4-yl.

The particularly preferred embodiments of the intermediates with respect to the variables correspond to those of the radicals R¹, R², R³ and R⁴ of formula I.

With respect to their intended use, preference is given to pyrimidines of formula I having the following substituents, where the preference is valid in each case on its own or in combination:

Compounds of formula I are preferred wherein R¹ denotes C₃-C₁₀-alkyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-C₁-C₆-alkyl, C₁-C₁₀-haloalkyl or phenyl being optionally substituted by one to three halogen atoms or C₁-C₁₀-alkyl or C₁-C₁₀-alkoxy.

12

Furthermore, particular preference is given to compounds I in which R¹ is C₁-C₁₀-haloalkyl, preferably polyfluorinated alkyl, in particular 2,2,2-trifluoroethyl, 2-(1,1,1-trifluoropropyl) or 2-(1,1,1-trifluorobutyl).


5

Likewise, particular preference is given to compounds I in which R¹ denotes optionally substituted C₃-C₈-cycloalkyl, preferably cyclopentyl or cyclohexyl.

- 10 Moreover, particular preference is given to compounds I in which R² represents phenyl being substituted by 2 or 3 substituents. Most preferred at least one of these substituents is attached in the 2-position with respect to the point of attachment to the pyrimidine moiety. Such substituents preferably include halogen
15 or alkoxy.

Furthermore, particular preference is given to compounds I in which R² represents a phenyl group of formula

20



- 25 wherein L¹ through L⁴ each independently represent hydrogen, fluorine, chlorine or methoxy, in particular L¹ represents fluorine or chlorine, L² and L⁴ each independently represent hydrogen, fluorine or chlorine, and L³ represents hydrogen, fluorine, chlorine or methoxy.
- 30 Particular preference is given to compounds of formula I in which R³ is chlorine.

- Besides, particular preference is given to compounds I in which R⁴
35 represents hydrogen, C₁-C₆-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl; or phenyl-C₁-C₄-alkyl, wherein the phenyl ring may be substituted by one or two halogen atoms.

- Likewise, particular preference is given to compounds I in which
40 R⁴ is hydrogen, C₁-C₆-alkyl or benzyl, especially C₁-C₆-alkyl.

Particular preference is given to compounds I in which X is NR⁵ and R⁵ is hydrogen, C₁-C₁₀-alkyl or C₁-C₁₀-haloalkyl, in particular hydrogen.


45

Besides, particular preference is given to compounds I in which R⁵ represents C₁-C₆-alkyl, especially hydrogen or methyl.

13

Particular preference is also given to compounds I in which X represent NR^5 and R^1 together with the interjacent nitrogen atom form an optionally substituted heterocyclic ring, preferably an optionally substituted 3- to 7-membered heterocyclic ring, in particular a pyrrolidine, piperidine, tetrahydropyridine, in particular 1,2,3,6-tetrahydropyridine or azepane ring which is optionally substituted by one or more C_1 - C_{10} -alkyl groups.

Most preferred are the compounds of formula IA



IA

in which R^1 to R^5 have the meaning given in formula I, L^1 is F or Cl, L^2 and L^4 each independently are H, F or Cl, and L^3 is H, F, Cl or OCH_3 .

Likewise, most preferred are the compounds wherein R^3 is chlorine, X is NH, R^4 is C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl or C_2 - C_6 -alkynyl, especially C_1 - C_6 -alkyl, R^2 represents phenyl optionally substituted by one or more fluorine and/or chlorine atoms and/or methoxy groups.

Particularly preferred are following compounds of formula IA:

	R^1	R^5	R^3	R^4	L^1	L^2	L^3	L^4
	$\text{CH}(\text{CH}_3)\text{CF}_3$	H	Cl	CH_3	F	H	F	F
	$\text{CH}(\text{CH}_3)\text{CF}_3$	H	Cl	$(\text{CH}_2)_3\text{CH}_3$	F	H	F	F
30	$\text{CH}(\text{CH}_3)_2$	H	Cl	CH_3	F	H	F	F
	$\text{c-C}_3\text{H}_5$	H	Cl	CH_3	F	F	H	F
	azepan-1-yl	H	Cl	CH_3	H	F	H	F
	$(\text{CH}_2)_2\text{CH}(\text{CH}_3)(\text{CH}_2)_2$		Cl	CH_3	F	H	H	Cl
35	$\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$	H	Cl	CH_3	F	H	H	Cl
	$(\text{CH}_2)_2\text{CH}(\text{CH}_3)(\text{CH}_2)_2$		Cl	CH_3	F	H	F	F
	$\text{CH}(\text{CH}_3)\text{CF}_3$	H	Cl	CH_3	F	H	F	F

Included in the scope of the present Invention are (R) and (S) isomers of compounds of general formula I having a chiral center and the racemates thereof, and salts, N-Oxides and acid addition compounds.

The compounds according to formula I are superior through their valuable fungicidal properties, in particular their enhanced systemicity. For example, they can be used in agriculture or related fields for the control of phytopathogenic fungi such as

Alternaria solani, *Botrytis cinerea*, *Cercospora beticola*, *Cladosporium herbarum*, *Corticium rolfsi*, *Erysiphe graminis*, *Helminthosporium tritici repentis*, *Lepfosphaeria nodorum*, *Micronectriella nivalis*, *Monilinia fructigena*, *Mycosphaerella ligulicola*, *Mycosphaerella pinodes*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Uncinula necator* and *Venturia inaequalis*, in particular *Pyricularia oryzae*, *Rhizoctonia solani* and *Venturia inaequalis*. The compounds of formula I according to the invention possess a high fungicidal activity within a wide concentration range.

Due to excellent activity, the compounds of formula I may be used in cultivation of all plants where infection by phytopathogenic fungi is not desired, e.g. cereals, solanaceous crops, vegetables, legumes, apples, vine.

The Invention further provides a fungicidal composition which comprises an active ingredient, which is at least one compound of formula I as defined above, and one or more carriers. A method of making such a composition is also provided which comprises bringing a compound of formula I as defined above into association with the carrier(s). Such a composition may contain a single active ingredient or a mixture of several active ingredients of the present Invention. It is also envisaged that different isomers or mixtures of isomers may have different levels or spectra of activity and thus compositions may comprise individual isomers or mixtures of isomers.

A composition according to the Invention preferably contains from 0.5% to 95% by weight (w/w) of active ingredient.

A carrier in a composition according to the Invention is any material with which the active ingredient is formulated to facilitate application to the locus to be treated, which may for example be a plant, seed, soil, or water in which a plant grows, or to facilitate storage, transport or handling. A carrier may be a solid or a liquid, including material which is normally a gas but which has been compressed to form a liquid.

The compositions may be manufactured into e.g. emulsion concentrates, solutions, oil in water emulsions, wettable powders, soluble powders, suspension concentrates, dusts, granules, water dispersible granules, micro-capsules, gels, tablets and other formulation types by well-established procedures. These procedures include intensive mixing and/or milling of the active ingredients with other substances, such as fillers, solvents, solid carriers, surface active compounds (surfactants), and optionally solid and/

15

or liquid auxiliaries and/or adjuvants. The form of application such as spraying, atomizing, dispersing or pouring may be Chosen like the compositions according to the desired objectives and the given circumstances.

5

Solvents may be aromatic hydrocarbons, e.g. Solvess® 200, substituted naphthalenes, phthalic acid esters, such as dibutyl or dioctyl phthalate, aliphatic hydrocarbons, e.g. cyclohexane or Paraffins, alcohols and glycols as well as their ethers and
10 esters, e.g. ethanol, ethyleneglycol mono- and dimethyl ether, ketones such as cyclohexanone, strongly polar solvents such as N-methyl-2-pyrrolidone, or γ -butyrolactone, higher alkyl pyrrolidones, e.g. n-octylpyrrolidone or cyclohexylpyrrolidone, epoxidized plant oil esters, e.g. methylated coconut or soybean oil ester
15 and watet. Mixtures of different liquids are offen suitable.

Solid carriers, which may be used for dusts, wettable powders, watet dispersible granules, or granules, may be mineral fillers, such as calcite, talc, kaolin, montmorillonite or attapulgate.

20 The physical properties may be improved by addition of highly dispersed silica gel or Polymers. Carriers for granules may be porous material, e.g. pumice, kaolin, sepiolite, bentonite; non-sorptive carriers may be calcite or sand. Additionally, a multitude of pregranulated inorganic or organic materials may be used,
25 such as dolomite or crushed plant residues.

Pesticidal compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application. The presence of small amounts of a carrier
30 which is a surfactant facilitates this process of dilution. Thus, preferably at least one carrier in a composition according to the Invention is a surfactant. For example, the composition may contain at two or more carriers, at least one of which is a surfactant.

35

Surfactants may be nonionic, anionic, cationic or zwitterionic substances with good dispersing, emulsifying and wetting properties depending an the nature of the compound according to general formula I to be formulated. Surfactants may also mean
40 mixtures of individual surfactants.

The compositions of the Invention may for example be formulated as wettable powders, water dispersible granules, dusts, granules, tablets, solutions, emulsifiable concentrates, emulsions,
45 suspension concentrates and aerosols. Wettable powders usually contain 5 to 90% w/w of active ingredient and usually contain in addition to solid inert carrier, 3 to 10% w/w of dispersing and

wetting agents and, where necessary, 0 to 10% w/w of stabilizer(s) and/or other additives such as penetrants or stickers. Dusts are usually formulated as a dust concentrate having a similar composition to that of a wettable powder but without a dispersant, and may be diluted in the field with further solid carrier to give a composition usually containing 0.5 to 10% w/w of active ingredient. Water dispersible granules and granules are usually prepared to have a size between 0.15 mm and 2.0 mm and may be manufactured by a variety of techniques. Generally, these types of granules will contain 0.5 to 90% w/w active ingredient and 0 to 20% w/w of additives such as stabilizer, surfactants, slow release modifiers and binding agents. The so-called "dry flowables" consist of relatively small granules having a relatively high concentration of active ingredient. Emulsifiable concentrates usually contain, in addition to a solvent or a mixture of solvents, 1 to 80% w/v active ingredient, 2 to 20% w/v emulsifiers and 0 to 20% w/v of other additives such as stabilizers, penetrants and corrosion inhibitors. Suspension concentrates are usually milled so as to obtain a stable, nonsedimenting flowable product and usually contain 5 to 75% w/v active ingredient, 0.5 to 15% w/v of dispersing agents, 0.1 to 10% w/v of suspending agents such as protective colloids and thixotropic agents, 0 to 10% w/v of other additives such as defoamers, corrosion inhibitors, stabilizers, penetrants and stickers, and water or an organic liquid in which the active ingredient is substantially insoluble; certain organic solids or inorganic salts may be present dissolved in the formulation to assist in preventing sedimentation and crystallization or as antifreeze agents for water. Aqueous dispersions and emulsions, for example compositions obtained by diluting the formulated product according to the invention with water, also lie within the scope of the invention.

Of particular interest in enhancing the duration of the protective activity of the compounds of this invention is the use of a carrier which will provide slow release of the pesticidal compounds into the environment of a plant which is to be protected.

The biological activity of the active ingredient can also be increased by including an adjuvant in the spray dilution. An adjuvant is defined here as a substance which can increase the biological activity of an active ingredient but is not itself significantly biologically active. The adjuvant can hitherto be included in the formulation as a coformulant or carrier, or can be added to the spray tank together with the formulation containing the active ingredient.

17

As a commodity the compositions may preferably be in a concentrated form whereas the end user generally employs diluted compositions. The compositions may be diluted to a concentration down to 0.001 % of active ingredient. The doses usually are in the range 5 from 0.01 to 10 kg a.i./ha.

Examples of formulations according to the Invention are:

Emulsion Concentrate (EC)

10	Active Ingredient	Compound of Example 5	30 % (w/v)
	Emulsifier(s)	Atlox® 4856 B/Atlox® 4858 B ¹⁾ (mixture containing calcium alkyl aryl sulfonate, fatty alcohol ethoxylates and light aromatics/mixture containing calcium alkyl aryl sulfonate, fatty alcohol ethoxylates and light aromatics)	5 % (w/v)
15			
20	Solvent	Shellsol® A ²⁾ (mixture of C ₉ -C ₁₀ -aromatic hydrocarbons)	to 1000 ml

Suspension Concentrate (SC)

25	Active Ingredient	Compound of Example 5	50 % (w/v)
	Dispersing agent	Soprophor® FL ³⁾ (polyoxyethylene polyaryl phenyl ether phosphate amine salt)	3 % (w/v)
30	Antifoaming agent	Rhodorsil® 422 ³⁾ (nonionic aqueous emulsion of polydimethylsiloxanes)	0.2 (w/v)
	Structure agent	Kelzan® S ⁴⁾ (Xanthan gum)	0.2 (w/v)
35	Antifreezing agent	Propylene glycol	5 % (w/v)
	Biocidal agent	Proxel® ⁵⁾ (aqueous dipropylene glycol solution containing 20% 1,2-beniso-thiazolin-3-one)	0.1 (w/v)
40			
	Water		to 1000 ml

18

Wettable Powder (WP)

Active Ingredient	Compound of Example 7	60 % (w/w)
Wetting agent	Atlox® 4995 ¹⁾	2 % (w/w)
5	(polyoxyethylene alkyl ether)	
Dispersing agent	Witcosperse® D-60 ⁶⁾	3 % (w/w)
	(mixture of sodium salts of condensed	
	naphthalene sulfonic acid and	
10	alkylarylpolyoxy acetates	
Carrier / Filter	Kaolin	35 % (w/w)

Water Dispersible Granules (WG)

15	Active Ingredient	Compound of Example 7	50 % (w/w)
	Dispersing /	Witcosperse® D-450 ⁶⁾	8 % (w/w)
	Binding agent	(mixture of sodium salts of condensed	
		naphthalene sulfonic acid and	
20		alkyl sulfonates)	
	Wetting agent	Morwet® EFW ⁶⁾	2 % (w/w)
		(formaldehyde condensation product)	
		Antifoaming agent	
		Rhodorsil® EP 6703 ³⁾	1 % (w/w)
25		(encapsulated silicone)	
	Disintegrant	Agrimer® ATF')	2 % (w/w)
		(cross-linked homopolymer of N-vinyl-2	
		pyrrolidone)	
	Carrier & Filter	Kaolin	35 % (w/w)

30

- 1) commercially available from ICI Surfactants
- 2) commercially available from Deutsche Shell AG
- 3) commercially available from Rhône-Poulenc
- 4) commercially available from Kelco Co.
- 35 5) commercially available from Zeneca
- 6) commercially available from Witco
- 7) commercially available from International Speciality Products

The compositions of this invention can be applied to the plants
 40 or their environment simultaneous with or in succession with
 other active substances. These other active substances can be Bi-
 ther fertilisers, agents which donate trace elements or other
 preparations which influence plant growth. However, they can also
 be selective herbicides, insecticides, fungicides, bactericides,
 45 nematicides, algicides, molluscicides, rodenticides, virucides,
 compounds inducing resistance into plants, biological control
 agents such as viruses, bacteria, nematodes, fungi and other mi-

croorganisms, repellents of birds and animals, and plant growth regulators, or mixtures of several of these preparations, if appropriate together with other carrier substances conventionally used in the art of formulation, surfactants or other additives
5 which promote application.

Furthermore, the other pesticide can have a synergistic effect anthe pesticidal activity of the compound of formula I.

10 The other fungicidal compound can be, for example, one which is also capable of combating diseases of cereals (e.g. wheat) such as those caused by *Erysiphe*, *Puccinia*, *Septoria*, *Gibberella* and *Helminthosporium* spp., seed and soil Borne diseases and downy and powdery mildews an eines, early and late blight an solanaceous
15 crops, and powdery mildew and scab an apples etc. These mixtures of fungicides can have a broader spectrum of activity than the compound of general formula I alone. Furthermore, the other fungicide can have a synergistic effect an the fungicidal activities of the compound of formula I.

20 Examples of the other fungicidal compounds are anilazine, azoxystrobin, benalaxyl, benomyl, binapacryl, bitertanol, blastocidin S, Bordeaux mixture, bromuconazole, bupirimate, captafol, captan, carbendazim, carboxin, carpropamid, chlorbenzthiazon, chlorothalonil, chlozolinat, copper-containing compounds such as copper
25 oxychloride, and copper sulfate, cycloheximide, cymoxanil, cyprofuram, cyproconazole, cyprodinil, dichlofluanid, dichlone, dichloran, diclobutrazol, diclocymet, diclomezine, diethofencarb, difenoconazole, diflumetorim, dimethirimol, dimethomorph, diniconazole, dinocap, ditalimfos, dithianon, dodemorph, dodine, edifenphos, epoxiconazole, etaconazole, ethirimol, etridiazole, famoxadone, fenapanil, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fenoxanil, fencpiclonil, fenpropidin, fenpropimorph, fentin, fentin acetate, fentin hydroxide, ferim-
30 zone, fluazinam, fludioxonil, flumetover, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, fosetylaluminium, fuberidazole, furalaxyl, furametpyr, guazatine, hexaconazole, IKF-916, imazalil, iminoctadine, ipconazole, iprodione, isoprothiolane, iprovalicarb, kasugamycin, KH-7281, kitazin P,
40 kresoximmethyl, mancozeb, maneb, mepanipyrim, mepronil, metalaxyl, metconazole, methfuroxam, MON 65500, myclobutanil, neoazoxin, nicket dimethyldithiocarbamate, nitrothalisopropyl, nuarimol, ofurace, organo mercury compounds, oxadixyl, oxycarboxin, penconazole, pencycuron, phenazineoxide, phthalide, polyoxin D,
45 polgram, probenazole, prochloraz, procymidione, propamocarb, propiconazole, propineb, pyraclostrobin, pyrazophos, pyrifenoxy, pyrimethanil, pyroquilon, pyroxyfur, quinomethionate, quinoxifen,

quintozene, spiroxamine, SSF-126, SSF-129, streptomycin, sulfur, tebuconazole, tecloftalame, tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tolclofosmethyl, tolylfluanid, triadimefon, triadimenol, triazbutil, triazoxide, 5 tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin A, vinclozolin, XRD-563, zarilamid, zineb, ziram.

In addition, the co-formulations according to the Invention may 10 contain at least one compound of formula I and any of the following classes of biological control agents such as viruses, bacteria, nematodes, fungi, and other microorganisms which are suitable to control insects, weeds or plant diseases or to induce host resistance in the plants. Examples of such biological control 15 agents are: *Bacillus thuringiensis*, *Verticillium lecanii*, *Autographa californica* NPV, *Beauveria bassiana*, *Ampelomyces quisqualis*, *Bacillus subtilis*, *Pseudomonas chlororaphis*, *Pseudomonas fluorescens*, *Steptomyces griseoviridis* and *Trichoderma harzianum*.

20 Moreover, the co-formulations according to the Invention may contain at least one compound of formula I and a chemical agent that induces the systemic acquired resistance in plants such as for example isonicotinic acid or derivatives thereof, 2,2-dichloro-3,3-dimethylcyclopropanecarboxylic acid or BION.

25 The compounds of formula I can be mixed with soil, peat or other rooting media for the protection of the plants against seed-borne, soilborne or foliar fungal diseases.

30 The Invention still further provides the use as a fungicide of a compound of the formula I as defined above or a composition as defined above, and a method for combating fungus at a locus, which comprises treating the locus, which may be for example plants subject to or subjected to fungal attack, seeds of such 35 plants or the medium in which such plants are growing or are to be grown, with such a compound or composition.

The present Invention is of wide applicability in the protection of crop and ornamental plants against fungal attack. Typical 40 crops which may be protected include cereals, grain crops such as wheat and barley, rice, sugar Beet, top fruit, peanuts, potatoes, vegetables and tomatoes. The duration of the protection is normally dependent on the individual compound selected, and also a variety of external factors, such as climate, whose impact is 45 normally mitigated by the use of a suitable formulation.

Synthesis Examples


With due modification of the starting compounds, the protocols shown in the synthesis examples below were used for obtaining 5 further compounds I. The resulting compounds, together with physical data, are listed in the Tables which follow.

Example 1

- 10 Preparation of 2-(N-cyano-N-methylamino)-4-chloro-5-(2,4,6-trifluorophenyl)-6-(1,1,1-trifluoroprop-2-ylamino)-pyrimidine

A mixture of 5-chloro-6-(2,4,6-trifluorophenyl)-7-(1,1,1-trifluoroprop-2-ylamino)-triazolo[1.5a]pyrimidine (2.5 g, 6.3 mmol, prepared according to WO-A 98/46608); dimethylformamide (15ml), sodium hydride (0.25 g, 60 %) and methyl iodide is stirred at ambient temperature for 45 minutes. The reaction mixture is poured into water (400 ml) and extracted with diethylether twice (300 ml). The organic Phase is separated, dried with anhydrous sodium sulphate and filtered. The filtrate is evaporated under reduced Pressure and purified by flash chromatography to yield 0.2 g of the product as a colorless oil.

Examples 2-20: Table I (synthesized analogously to Ex. 1)



30

Ex.	R ¹	R ⁵	R ⁴	L ¹	L ²	L ³	L ⁴	melting point (°C)	
35	2	1,1,1-trifluoro-prop-2-yl	H	butyl	F	H	F	F	168-172
	3	iso-propyl	H	methyl	F	H	F	F	130
	4	cyclopentyl	H	methyl	F	F	H	F	140
	5	-(CH ₂) ₂ -CH(CH ₃)-(CH ₂) ₂ -		methyl	F	H	H	Cl	oil
40	6	2-butyl	H	methyl	F	H	H	Cl	oil
	7	-(CH ₂) ₂ -CH(CH ₃)-(CH ₂) ₂ -		methyl	F	H	F	F	100
	8	1,1,1-trifluoro-prop-2-yl	methyl	methyl	F	H	F	F	86
45	9	but-2-yl	ethyl	methyl	F	H	F	F	
	10	norborn-2-yl	H	methyl	F	H	F	F	
	11	-(CH ₂) ₂ -CH(CH ₃)-(CH ₂) ₂ -		ethyl	F	H	H	Cl	

22

Ex.	R ¹	R ⁵	R ⁴	L ¹	L ²	L ³	L ⁴	melting point (°C)
5	12	cyclopentyl	H	methyl	F	H	F	F
	13	iso-propyl	H	methyl	F	H	H	Cl
	14	ethyl	ethyl	methyl	F	H	F	F
10	15	2,2,2-trifluoro-ethyl	H	methyl	F	H	F	F
	16	1,1,1-trifluoro-prop-2-yl	H	ethyl	F		H	Cl
	17	2-butyl	H	methyl	F	H	H	Cl
15	18	norborn-2-yl	H	methyl	F	H	H	Cl
	19	1,1,1-trifluoro-prop-2-yl	H	methyl	F	H	OCH ₃	F
	20	methallyl	ethyl	methyl	F	H	F	F
20	21	2,2,2-trifluoro-ethyl	H	allyl	F	H	H	Cl
	22	1,1,1-trifluoro-prop-2-yl	H	n-propyl	F	H	F	F
	23	-(CH ₂) ₇ -		methyl	H	F	H	F
	24	iso-propyl	methyl	methyl	F	H	H	Cl
	25	1,1,1-trifluoro-prop-2-yl	H	methyl	F	H	F	F

25

Example 26

Preparation of

2-(N-cyano-N-methylamino)-4-chloro-5-(2,4,6-trifluorophenyl)-6-cyclohexylpyrimidine


A mixture of 5-chloro-6-(2,4,6-trifluorophenyl)-7-cyclohexyltriazolo[1.5a]pyrimidine (2.5 g, 6.3 mmol, prepared according to WO-A 99/41255), dimethylformamide (15ml), sodium hydride (0.25 g, 60 %) and methyl iodide is stirred at ambient temperature for 45 minutes. The reaction mixture is poured into water (400 ml) and extracted with diethylether twice (300 ml). The organic phase is separated, dried with anhydrous sodium sulphate and filtered. The filtrate is evaporated under reduced pressure and purified by flash chromatography to yield 0.2 g of the product as a colorless oil.

45

23

Examples 27-39: Table II (synthesized analogously to Ex. 26)

5



Ex.	R ¹	L ¹	L ³	L ⁴	melting point (°C)
27	n-heptyl	F	F	F	
28	cyclopentyl	F	F	F	
29	cyclohexyl	F	F	F	
30	4-methylcyclohexyl	F	F	F	
31	2-methylpropyl	F	F	F	
32	n-heptyl	F	H	Cl	
33	cyclopentyl	F	H	Cl	
34	cyclohexyl	F	H	Cl	
35	n-hexyl	F	H	Cl	
36	4-methylcyclohexyl	F	H	Cl	
37	2-methylpropyl	F	H	Cl	
38	4-fluorocyclohexyl	F	F	F	
39	4-fluorocyclohexyl	F	OCH ₃	F	

Example 40

30 Preparation of 4-chloro-2-(N-cyanoamino)-6-[(4-methyl)-piperidin-1-yl]-5-phenyl-pyrimidine

To a solution of 4-chloro-6-[(4-methyl)-piperidin-1-yl]-2-methyl-sulfony-5-phenyl-pyrimidine (1.0g, 2.7 mmol) in dimethylformamide (8ml) at room temperature was added potassium carbonate (0.76g, 5.47mmol). After the reaction mixture had been stirred at room temperature for 17.5 hours the reaction mixture was then diluted with water (70ml) and the resulting cloudy solution acidified to pH 1 by the addition of concentrated hydrochloric acid (4ml). The resulting white suspension was then stirred at room temperature for 2 hours. The suspension was then filtered, washed with water followed by hexane and dried in vacuo overnight. The crude product was recrystallized from methylene chloride/hexane to afford 0.71 g (79% yield) of the title compound as a white crystalline solid (melting point: 219-220°C (dec)).

Example 41

Preparation of

4-chloro-2-(N-cyano-N-methylamino)-6-[(4-methyl)-piperi-
5 din-1-yl]-5-phenylpyrimidine

To a solution of 4-chloro-2-(N-cyanoamino)-6-[(4-methyl)-pi-
peridin-1-yl]-5-phenylpyrimidine (0.1 g, 0.305mmol) in dimethyl-
formamide (4ml) at room temperature was added water (2ml) follo-
10 wed by potassium carbonate (0.084g, 0.61 mmol) and the resulting
suspension was warmed gently to afford a clear solution. To the
cooled solution was then added methyl iodide (0.076ml, 1.22mmol)
and the reaction mixture stirred at room temperature for 2.5
hours. The reaction was then quenched by the addition of satura-
15 ted aqueous ammonium chloride solution (40ml). After adding ethyl
acetate (40ml) the biphasic mixture was stirred for 5 minutes.
The organic phase was then separated, washed with saturated brine
(50ml), dried over magnesium sulfate and concentrated in vacuo to
afford a yellow syrup. The crude product was chromatographed an
20 silica gel eluting with 90:10 v/v hexane: ethyl acetate to afford
0.09g (87% yield) of the title compound as a colorless syrup.

Example 42

25 Preparation of

2-(N-benzyl-N-cyano)-4-chloro-6-[(4-methyl)-piperi-
din-1-yl]-5-phenylpyrimidine

To a solution of 4-chloro-2-(N-cyanoamino)-6-[(4-methyl)-piperi-
30 din-1-yl]-5-phenylpyrimidine (0.37g, 1.13mmol) in dimethylform-
amide (10ml) at room temperature was added water (3ml) and potas-
sium carbonate (0.19g, 1.35mmol). To the resulting milky suspen-
sion benzyl bromide (0.16ml, 1.35mmol) was added and stirred at
room temperature for 18.75 hours before being quenched by the ad-
35 dition of a saturated aqueous ammonium chloride solution (40ml).
The mixture was then partitioned between ethyl acetate (75ml) and
water (75ml), dried over magnesium sulfate and concentrated in
vacuo to afford a yellow syrup. The crude product was chromato-
graphed on silica gel eluting with 90:10 v/v hexane: ethyl ace-
40 tate to afford 0.47g (100% yield) of the title compound as a
white crystalline solid (melting point 98-100°C).

Example 43

Alternative preparation of
4-chloro-2-(N-cyano-N-methylamino)-6-[(4methyl)-piperi-
5 din-1-yl]-5-phenylpyrimidine

To a solution of 4-chloro-6-[(4-methyl)-piperidin-1-yl]-2-methyl-
sulfonyl-5-phenyl-pyrimidine (0.5g, 1.37mmol) in dimethylform-
amide (6ml) was added potassium carbonate (0.38g, 2.73mmol) fol-
10 lowed by methyl cyanamide (0.31 g, 5.47mmol). After stirring the
reaction mixture at room temperature for 19 hours the reaction
mixture was diluted with water (75ml), and extracted with ethyl
acetate (75ml). The organic phase was washed with water (75ml),
followed by saturated brine (75ml), dried over magnesium sulfate
15 and concentrated in vacuo to afford a yellow syrup. The crude
product was chromatographed on silica gel eluting with 90:10 v/v
hexane: ethyl acetate to afford 0.39g (84% yield) of the title
compound as a colorless syrup.

20 Example 44

Preparation of the starting material 4-chloro-6-[(4-methyl)-pi-
peridin-1-yl]-2-methylsulfonyl-5-phenylpyrimidine

25 Step a: 5-Phenyl-2-methylthio-4,6(1H,5H)-pyrimidinedione

60.0 g (208 mmol) of ethyl 2-phenylmalonate and 19.0 g (249 mmol)
of thiourea were heated at 150°C for 2.5 hours in 77 g (416 mmol)
of tri-n-butylamine. The resultant ethanol was for the most part
30 distilled off. After the reaction solution had cooled, 180 ml of
an aqueous solution of 24.9 g (623 mmol) of NaOH were added to
it. After adding 50 ml of cyclohexane and stirring for about
30 min the aqueous phase was separated off, treated with 35.4 g
(142 mmol) of methyl iodide and stirred at approximately 20 to
35 25°C for about 16 h. After acidification with dilute HCl solution
and stirring for about 30 min the precipitate was filtered off.
After washing with water and drying, 16.7 g of the compound in
the title (28 % of theoretical) were obtained in the form of
white crystals.

40

Step b: 4,6-Dichloro-5-phenyl-2-methylthiopyrimidine

A solution of 48.8 g (170 mmol) of the product from step A in
200 ml of phosphoryl chloride to which 3 ml of dimethylformamide
45 (DMF) had been added was refluxed for 40 hours. After distilling
off most of the phosphoryl chloride and diluting the residue with
ethyl acetate, water was added while stirring at 15 to 20°C. After

phase separation, the organic phase was washed with water and dilute NaHCO_3 solution, then dried and freed of solvent. 37.5 g of the compound in the title (68 % of theoretical) were obtained in the form of an oil which was employed in step C without further purification.

Step c: 6-Chloro-5-phenyl-4-[(4-methyl)piperidin-1-yl]-2-methylthiopyrimidine

- 10 A solution of 37.5 g (324 mmol) of the product from step B in 150 ml of anhydrous dichloromethane was treated with 24 g (406 mmol) of isopropylamine and stirred for five hours at about 20 to 25°C. The solvent was then distilled off, the residue taken up in ethyl acetate and washed with dilute HCl, water and dilute NaHCO_3 solution, then dried and freed of solvent. After chromatographing the residue on silica gel (cyclohexane/methyl tert-butyl ether 100:1 to 19:1) 13.4 g of the title compound (33 % of theoretical) were obtained in the form of colorless crystals which were employed in the next stage without further purification.

20

Step d: Preparation of 6-chloro-4-[(4-methyl)-piperidin-1-yl]-2-methylsulfonyl-5 phenylpyrimidine

- A solution of 4-chloro-6-[(4-methyl)-piperidin-1-yl]-2-methylthio-5 phenylpyrimidine (17.19g, 51.5mmol) in methylene chloride (350ml) was cooled to 0°C. After adding 70% in chloroperbenzoic acid (25.4g, 103mmol) the reaction mixture was stirred at 0°C for 1 hour and at room temperature for 1.5 hours. The reaction mixture was then concentrated in vacuo to a small volume, diluted with ethyl acetate (400 ml), washed with 5% aqueous solution of sodium carbonate (3 x 300ml) and saturated brine (300ml), dried over magnesium sulfate and concentrated in vacuo to afford a white solid which was then recrystallized from ethyl acetate/hexane to afford 14.68g (78% yield) of the title compound as an off-white crystalline solid (melting point: 160-162°C).

Example 45

Preparation of the intermediate

- 40 4-chloro-6-(N-cyclopentyl)amino-5-(2-fluorophenyl)-2-methylthiopyrimidine

Step a: Preparation of 5-(2-fluorophenyl)-2-thiobarbituric acid

- 45 To absolute ethanol (200ml), sodium (3.62g, 157mmol) was added at room temperature under nitrogen atmosphere. The reaction mixture was stirred until all the sodium had reacted. A solution of

27

diethyl-(2-fluorophenyl)malonate (20g, 78.7mmol) in absolute ethanol (50ml) was then added followed by thiourea (8.38g, 110mmol). The reaction mixture was then refluxed under nitrogen atmosphere for 17 hours. The cooled reaction mixture was then poured into water (800ml), the reswiting mixture stirred for 15 minutes and extracted with diethyl ether (500ml). The organic phase was extracted with brine (150ml, 1/3 saturated) and the aqueous layers combined. Now concentrated hydrochloric acid (14ml) was added to the aqueous phase and the resulting white suspension was stirred gently for 1 hour. The suspension was filtered and the resulting solid washed with water, followed by diethyl ether and dried in vacuo for approximately 12 hours to afford 8.59g (46% yield) of the title compound as a white solid (melting point: >240°C).

15

Step b: Preparation of

4,6-dihydroxy-5-(2-fluorophenyl)-2-methylthiopyrimidine

To a mixture of 5-(2-fluorophenyl)-2-thiobarbituric acid (8.20g, 34.3mmol) and a 2.0M aqueous solution of sodium hydroxide (68.8ml, 13.8mmol) was added dropwise over 30 minutes dimethyl sulfate (4.348, 34.4mmol) at room temperature with stirring. The reaction mixture was stirred then for an additional 24 hours at room temperature. After washing with ethyl acetate (2 x 100ml) the aqueous phase was acidified to pH 1 by adding concentrated hydrochloric acid (8ml). The resulting white suspension was stirred for 30 minutes, filtered and the resulting white solid washed with water followed by hexane and dried under vacuum to afford 7.11 g of a white solid. Ethyl acetate (100ml) was added to the crude product and the resulting suspension refluxed while stirring for 15 minutes. The cooled suspension was then filtered and dried to afford 5.41 g (61% yield) of the title compound as a white solid (m. p. > 240°C).

35 Step c: Preparation of

4,6-dichloro-5-(2-fluorophenyl)-2-methylthiopyrimidine

To a suspension of 4,6-dihydroxy-5-(2-fluorophenyl)-2-methylthiopyrimidine (0.75g, 2.97mmol) in phosphorous oxychloride (7.5ml, 80.5mmol) was added tri-n-propylamine (1.24ml, 6.54mmol) at room temperature under nitrogen atmosphere. After refluxing the reaction mixture for 18 hours and cooling to room temperature the reaction mixture was concentrated in vacuo and the resulting dark brown residue was dissolved in a minimum amount of acetonitrile and added to water (75ml) under stirring. Ethyl acetate (75ml) was then added and the resulting biphasic mixture stirred vigorously for one hour. The organic phase was separated, washed with

28

2M aqueous hydrochloric acid (75ml), saturated aqueous sodium bicarbonate solution (2 x 75 ml) and saturated brine (75ml). After drying over magnesium sulfate the organic Phase was concentrated in vacuo to afford a light brown oil. The crude product
5 was chromatographed on silica gel eluting with 98:2 v/v hexane: ethyl acetate to afford 0.74g (86% yield) of the title compound as colorless crystals.

Step d: Preparation of 4-chloro-6-(N-cyclopentyl)amino-5-(2-fluorophenyl)-2-methylthiopyrimidine
10

To a solution of 4,6-dichloro-5-(2-fluorophenyl)-2-methylthiopyrimidine (0.74g, 256mmol) in methylene chloride (1 ml) at room temperature under nitrogen atmosphere was added cyclopentyl-amine
15 (1.01 ml, 10.24mmol) and the reaction mixture was stirred at room temperature for 15 hours. The reaction mixture was then diluted with 1:1 v/v diethyl ether ethyl acetate (75ml), washed with 1 M aqueous hydrochloric acid (2 x 75ml), saturated aqueous sodium bicarbonate solution (75ml) and saturated brine (75ml), dried
20 over magnesium sulfate and concentrated in vacuo to afford a colorless syrup. The crude product was chromatographed on silica gel eluting with 95:5 v/v hexane: ethyl acetate to afford 0.87g (100% yield) of the title compound as a white crystalline solid (melting point: 81-83°C).

25

Example 46

Preparation of the intermediate

4,6-dichloro-5-(2-chloro-6-fluorophenyl)-2-methylthiopyrimidine

30

Step a: Preparation of 4,6-dihydroxy-5-(2-chloro-5-fluorophenyl)-2-methylthiopyrimidine

A mixture of diethyl(2-chloro-6-fluorophenyl)malonate (12.01 g,
35 41.6mmol), thiourea (3.8g, 49.92mmol) and tributylamine (19.82ml, 83.2mmol) was stirred at 150°C under nitrogen atmosphere for 3 hours. The cooled reaction mixture was then partitioned between ethyl acetate (84ml) and 2.0M sodium hydroxide solution (83.2ml, 106mmol) with vigorous stirring for 15 minutes. The aqueous phase
40 was separated, dimethyl sulfate (5.258, 41.6mmol) added and the mixture stirred for approximately 12 hours at room temperature. Additional 5.0 sodium hydroxide solution (33.4ml, 166mmol) and dimethylsulfate (2.63g, 20.8mmol) were added and the mixture stirred for further 2 hours. The resulting suspension was then filtered,
45 the filtrate acidified to pH 1 by the addition of concentrated hydrochloric acid and stirred for 30 minutes. The resulting suspension was then filtered and the resulting white solid washed


with water followed by hexane and dried under vacuum approximately 12 hours to afford 2.57g (22%yield) of the title compound as a white solid.

5 Step b: Preparation of 4,6-dichloro-5-(2-chloro-6-fluorophenyl)-2-methylthiopyrimidine

To a suspension of 4,6-dihydroxy-5-(2-chloro-6-fluorophenyl)-2-methylthiopyrimidine (2.57g, 8.96mmol) in phosphorous
 10 oxychloride (25.7ml, 276mmol) was added tri-n-propylamine (3.75ml, 19.72mmol) at room temperature under nitrogen atmosphere. After refluxing the reaction mixture at 140°C for 40 hours and cooling to room temperature the reaction mixture was concentrated in vacuo to afford a black oil. The crude product was par-
 15 titioned between ethyl acetate (250ml) and water (250ml) and the resulting biphasic mixture stirred vigorously for 15 minutes. The organic phase was separated, washed with 2M aqueous hydrochloric acid (2 x 250ml), saturated sodium bicarbonate solution (2x 250ml) and saturated brine (250ml). After drying over magnesium
 20 sulfate the organic Phase was concentrated in vacuo to afford 2.3g (79% yield) of the title compound as a dark brown solid.

Examples 47-49: Table III (synthesized analogously to Ex. 40-46)

25



IA

30

Ex.	R ¹	R ⁵	R ⁴	melting point (°C)
47	CF ₃ CH(CH ₃)	H	C ₆ H ₅ -CH ₂	178
48	(S)-CF ₃ CH(CH ₃)	H	CH ₃ O-C(=O)C(=CH ₂)CH ₂	120-123
49	CF ₃ CH(CH ₃)	H	(CH ₃ CH ₂ O-C[=O]) ₂ C(CH ₃)	178
50	CF ₃ CH(CH ₃)	H	CH=C-CH ₂	185-186
51	CF ₃ CH(CH ₃)	H	thien-3-yl-CH ₂	68-72

40 Biological Investigations

A Determination of Minimum Inhibitory Concentration by Test Compounds in the Serial Dilution Test with *Pyricularia Oryzae*

45 The MIC (Minimum Inhibitory Concentration) value, which indicates the lowest concentration of the active ingredient in the growth medium which causes a total inhibition of mycelial growth, is

determined by serial dilution tests using Microtiter plates with 24 or 48 wells per plate. The dilution of the test compounds in the nutrient solution and the distribution to the wells is carried out by a TECAN RSP 5000 Robotic Sample Processor. The following test compound concentrations are used: 0.05, 0.10, 0.20, 0.39, 0.78, 1.56, 3.13, 6.25, 12.50, 25.00, 50.00 and 100.00 µg/ml. For preparation of the nutrient solution, V8 vegetable Juice (333 ml) is mixed with calcium carbonate (4.95 g), centrifuged, the supernatant (200 ml) diluted with water (800 ml) and autoclaved at 121 °C for 30 min.

The inocula of *Pyricularia Oryzae* are added into the wells as spore suspensions (50 µl; 5×10^5 /ml) or agar slices (6 mm) of an agar culture of the fungus.

After 6-12 days incubation at temperatures of 18 to 25°C, the MIC values are determined by visual inspection of the plates, as shown in Table IV.

Table IV

Example No.	MIC [µg/ml]
3	3.13
4	> 100
5	25
6	6.25
7	100

B Evaluation of In Vivo Fungicidal Activity of Test Compounds

Test compounds are dissolved in acetone and diluted with deionized water (95 parts water to 5 Parts acetone), containing 0.05% TWEEN 20®, a polyoxyethylene sorbitan monolaurate surfactant manufactured by Atlas Chemical Industries, to give a concentration of 200 ppm.

The plants are sprayed with the test solutions, dried and inoculated with fungi later the same day. When disease symptom development is optimal, the plants are rated for disease control according to the rating scale shown below. Each test contains inoculated treated plants, inoculated untreated plants and inoculated plants treated with reference fungicides. The data obtained are shown in Table V.

RATING SCALE

Rating	Range % Control
0	0
5 1	1-14
2	15-29
3	30-44
4	45-59
5	60-74
10 6	75-89
7	90-95
8	96-99
9	100

15 TARGETS

Symbol	Disease	Pathogen
AS	Apple Scab	<i>Venturia inaequalis</i>
RB	Rice Blast	<i>Pyricularia grisea</i> sp. <i>oryzae</i>
20 GDM	Grape downy mildew	<i>Uncinula necator</i>

Table V

	Example	AS	RB	GDM
25	2	0	0	0
	3	7	5	7
	4	5	3	3
	5	9	4	0
30	6	9	6	3

C Evaluation of In Vitro Fungicidal Activity against
Rhizoctonia solani

- 35 Test compounds are dissolved in acetone to give a concentration of 10 ppm and added to individual cell walls (24-cell-well plates, Corning), which were previously filled with a suspension of ground fungal mycelium in a chemically defined growth medium. After 3-7 days of incubation, inhibition of mycelial growth is
- 40 recorded using the following scale: The data obtained are shown in Table VI.

RATING SCALE

Rating	Degree of Inhibition
0	None
5 3	slight
5	moderate
7	severe
9	complete

10 Table VI

Example	<i>Rhizoctonia solani</i> (Rice sheath blight)
1	7
15 4	7
5	0
6	0
7	0
8	7
20	

25

30


35

40

45

Claims:

1. Pyrimidines of formula I



I

in which

R¹ represents hydrogen or

C₁-C₁₀-alkyl, C₁-C₁₀-haloalkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₄-C₈-alkadienyl, C₁-C₁₀-alkoxy, C₃-C₈-cycloalkyl, phenyl, or

5- or 6-membered heteroaryl or 5- or 6-membered heterocyclyl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, or

tri-C₁-C₆-alkyl-silyl, formyl or C₁-C₁₀-alkoxycarbonyl;

wherein R¹ groups are unsubstituted or substituted by one to three groups R^a

R^a halogen, nitro, cyano, hydroxy or

C₁-C₁₀-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₁-C₁₀-haloalkyl, C₃-C₆-halocycloalkyl, C₁-C₁₀-alkoxy, C₁-C₁₀-haloalkoxy, C₁-C₁₀-haloalkoxy, C₁-C₆-alkoxycarbonyl, tri-C₁-C₄-alkylsilyl, phenyl, halo- or dihalo-phenyl or 5- or 6-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom;

R² represents phenyl, C₃-C₆-cycloalkyl or 5- or 6-membered heteroaryl, containing one to four nitrogen atoms or one to three nitrogen atoms and one sulfur or oxygen atom, which are unsubstituted or substituted by one to three groups R^a;

R³ represents hydrogen, halogen or

34

C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy, C₁-C₁₀-alkylthio, C₁-C₁₀-alkylamino or di-C₁-C₁₀-alkylamino; which are unsubstituted or substituted by one to three groups R^a;

5 R⁴ represents hydrogen or

C₁-C₁₀-alkyl, C₂-C₆-alkenyl or C₂-C₆-alkynyl; which are unsubstituted or substituted by one to three groups R^a; and

10


X represents O, S, NR⁵ or a single bond, wherein R⁵ represents hydrogen or C₁-C₁₀-alkyl; or

15

R¹ and R⁵ together with the interjacent nitrogen atom form a heterocyclic ring.

2. Pyrimidines of formula I according to claim 1, in which R² represents a phenyl group of formula

20




25

wherein L¹ through L⁴ each independently represent hydrogen, fluorine, chlorine or methoxy.

3. Pyrimidines of formula IA

30



IA

35

in which R¹ to R⁵ have the meaning given in claim 1, and L¹ through L⁴ are as defined in claim 2.

4. Pyrimidines according to claim 1 to 3 in which R³ represents chlorine.


40

5. Pyrimidines according to claims 1 to 4 in which R⁴ represents hydrogen, C₁-C₆-alkyl or benzyl.

45

35

6. A process for the preparation of pyrimidines of formula I according to claim 1 wherein R⁴ is optionally substituted alkyl, alkenyl or alkynyl by treating compounds of the formula II



II


in which R¹ through R³ and X are as defined in formula I; with a base and an alkylation agent of formula III



III


in which R⁴ is C₁-C₆-alkyl, C₁-C₆-alkenyl or C₁-C₆-alkynyl; which are unsubstituted or substituted by one to three groups R^a, and Y represents halogen atom.

7. A process for the preparation of pyrimidines of formula I according to claim 1 wherein R⁴ is is C₁-C₆-alkyl, C₁-C₆-alkenyl or C₁-C₆-alkynyl which are unsubstituted or substituted by one to three groups R^a by reacting sulfones of formula VI




VI

in which R¹ through R³ and X are as defined in formula I and R⁶ is C₁-C₆-alkyl or C₁-C₆-haloalkyl; with alkylated cyanamides of formula VII



VII

in which R⁴ is is C₁-C₆-alkyl, C₁-C₆-alkenyl or C₁-C₆-alkynyl which are unsubstituted or substituted by one to three groups R^a; wherein sulfones of formula VI are obtained by reacting 2-thiopyrimidinederivatives of formula VIII



VIII

in which the variables are as defined in formula VI; with oxidizing agents.

8. Compounds of formulae VI and VIII as defined in claim 7.

5

9. A fungicidal composition which comprises a carrier and an effective amount of at least one compound of formula I as defined in claim 1.

10 10. A method of combating fungus at a locus which comprises treating the locus with an effective amount of at least one compound of formula I as defined in claim 1.

15

20

25

30

35

40

45

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06565

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/48 C07D401/04 A01N43/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 02513 A (FUJISAWA PHARMACEUTICAL CO) 20 February 1992 (1992-02-20) example 18	1,2,5
X	WO 85 00603 A (STERLING DRUG INC) 14 February 1985 (1985-02-14) page 14, line 20 - line 21 -/-	1,5



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *G* document member of the same patent family

Date of the actual completion of the international search

26 October 2001

Date of mailing of the international search report

13/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06565

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; BENNETT, GREGORY B. ET AL: "Synthesis and antiinflammatory activity of trisubstituted pyrimidines and triazines" retrieved from STN Database accession no. 89:36588 CA XP002181005 CAS RN 66521-94-6 abstract & J. MED. CHEM. (1978), 21(7), 623-8 ,	1,2,5
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; TAKAGI, KANAME ET AL: "Synthesis of pyrimidines and pyrazoles from 3-acyl-5-halobenzofurans" retrieved from STN Database accession no. 84:30999 CA XP002181006 CAS RNs 57832-35-6, 57832-36-7, 57832-37-8, 57832-38-9 abstract & CHEM. PHARM. BULL. (1975), 23(10), 2427-31 ,	1,2,5
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; TAKAGI, KANAME ET AL: "Antitumor agents. IV. Formation of new pyrimidines from benzofurans substituted in position 3 by an electron attracting group" retrieved from STN Database accession no. 74:3578 CA XP002181007 CAS RNs 29936-04-7, 29936-06-9, 29936-07-0, 30041-99-7 abstract & CHIM. THER. (1970), 5(4), 264-9 ,	1,2,5

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06565

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; GOMPPER, RUDOLF ET AL: "Synthesis of oligo(diazaphenyls). Tailor-made fluorescent heteroaromatics and pathways to nanostructures" retrieved from STN Database accession no. 127:176399 CA XP002181008 see CAS RNs attached to abstract & SYNTHESIS (1997), (6), 696-718 ,	8
X,P	WO 00 73278 A (HOFFMANN LA ROCHE) 7 December 2000 (2000-12-07) examples 73c, 80c	8
X	WO 98 38171 A (SIGNAL PHARM INC) 3 September 1998 (1998-09-03) examples 125,127,129	8
X	WO 97 09311 A (HOFFMANN LA ROCHE) 13 March 1997 (1997-03-13) examples 129(a),129(b)	8
X	WO 95 35293 A (VITA INVEST SA ;DEL CASTILLO NIETO JUAN CARLOS (ES); CALDERO GES J) 28 December 1995 (1995-12-28) example 30	8
X	WO 91 11441 A (HOECHST AG) 8 August 1991 (1991-08-08) example 15; table I, lines 20.35	8
X	WO 89 03416 A (MERCK PATENT GMBH) 20 April 1989 (1989-04-20) examples 17,18	8
X	EP 0 295 370 A (HOECHST AG) 21 December 1988 (1988-12-21) examples 1-7,10,12-24,33	8
X	EP 0 284 008 A (HOECHST AG) 28 September 1988 (1988-09-28) examples 1,4,6-9	8
X	EP 0 257 605 A (CIBA GEIGY AG) 2 March 1988 (1988-03-02) examples 1b), 1c)	8
X	US 4 438 117 A (CHERKOFKY SAUL C) 20 March 1984 (1984-03-20) examples 1(interm.(c)),2 (start.comp.),3,9,10,14	8
	-/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06565

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; PARK, JUNG HO ET AL: "A new synthesis of a chiral ester containing phenylpyrimidine ring as liquid crystal dopant" retrieved from STN Database accession no. 123:313859 CA XP002181009 see CAS RNs attached to abstract & BULL. KOREAN CHEM. SOC. (1995), 16(6), 489-92 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; FISCHER, GERHARD W.: "Tetrazole compounds. 8. Synthesis of tetrazolylpyrimidines from tetrazolyl-substituted enamino ketones" retrieved from STN Database accession no. 120:270297 CA XP002181010 see CAS RNs attached to abstract & J. HETEROCYCL. CHEM. (1993), 30(6), 1517-19 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; NOHIRA, HIROYUKI ET AL: "Optically active compounds as liquid crystal compositions for optical display devices" retrieved from STN Database accession no. 118:202602 CA XP002181011 see CAS RNs attached to abstract & JP 04 300871 A (CANON K. K., JAPAN) 23 October 1992 (1992-10-23)</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SANDOSHAM, JESSIE ET AL: "Stannylation reactions and palladium catalysis in the syntheses of unsymmetrical biheteroaryls" retrieved from STN Database accession no. 112:158185 CA XP002181012 see CAS RNs attached to abstract & ACTA CHEM. SCAND. (1989), 43(7), 684-9 ,</p>	8

-/-

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 01/06565

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; MAMAEV, V. P. ET AL: "Synthesis of 2,5'-bipyrimidines from substituted 5-cyanopyrimidines" retrieved from STN Database accession no. 111:134072 CA XP002181013 see CAS RN attached to abstract & KHIM. GETEROTSIKL. SOEDIN. (1988), (3), 371-5 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; MUELLER, INGRID ET AL: "New phenyl-pyrimidine derivatives with SC-phases" retrieved from STN Database accession no. 110:67361 CA XP002181014 see CAS RNs attached to abstract & FERROELECTRICS (1988), VOLUME DATE 1987, 85, 393-403 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SHARMA, N. K. ET AL: "Binary liquid crystal systems with two eutectics" retrieved from STN Database accession no. 108:159361 CA XP002181015 see CAS RNs attached to abstract & MOL. CRYST. LIQ. CRYST. (1987), 151(Pt. A), 225-31 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ZASCHKE, HORST ET AL: "2-Cyano-5-'4-acyloxyphenyl!-pyrimidines" retrieved from STN Database accession no. 95:97846 CA XP002181016 see CAS RNs attached to abstract & DD 145 913 Z (GER. DEM. REP.) 14 January 1981 (1981-01-14)</p>	8

-/-

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06565

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ZASCHKE, HORST ET AL: "Synthesis of low-melting-point, liquid-crystal heterocycles;III. 2-n-Alkylthio-5-(4-substituted-phenyl)pyri midines" retrieved from STN Database accession no. 87:184456 CA XP002181017 see CAS RNs attached to abstract & Z. CHEM. (1977), 17(8), 293-4 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ZASCHKE, HORST: "Liquid crystalline 2-cyanopyrimidines" retrieved from STN Database accession no. 87:5897 CA XP002181018 see CAS RNs attached to abstract & Z. CHEM. (1977), 17(2), 63-4 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; STREKOWSKI, LUCJAN: "Synthesis with pyrimidinyl lithium compounds and properties of a series of 4,4'- and 4,5'-bipyrimidine derivatives" retrieved from STN Database accession no. 86:189847 CA XP002181019 see CAS RN attached to abstract & SER. CHEM. - UNIW. ADAMA MICKIEWICZA POZNANIU (1976), 20, 114 PP. ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SRIVASTAVA, S. P. ET AL: "Identification of pyrimidines by thin-layer chromatography" retrieved from STN Database accession no. 85:201764 CA XP002181020 see CAS RN attached to abstract & FRESSENIUS' Z. ANAL. CHEM. (1976), 281(4), 302 ,</p>	8

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/06565

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; KRCHNAK, V. ET AL: "Synthetic reactions of dimethylformamide. XXIX. Synthesis of 2,2'-substituted 5,5'-bipyrimidines" retrieved from STN Database accession no. 82:72920 CA XP002181021 see CAS RN attached to abstract & COLLECT. CZECH. CHEM. COMMUN. (1974), 39(11), 3327-31 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; BROWN, DESMOND J. ET AL: "Dimroth rearrangement. XIII. The small effect of p-substitution on rearrangement rates for 1,2-dihydro-2-imino-1-methyl-5- phenylpyrimidines" retrieved from STN Database accession no. 74:75908 CA XP002181022 see CAS RNs attached to abstract & J. CHEM. SOC., C (1971), (2), 250-6 ,</p>	8
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; BROWN, DESMOND J. ET AL: "Simple pyrimidines. XII. Synthesis and methylation of 2-amino-5-phenylpyrimidines" retrieved from STN Database accession no. 74:112008 CA XP002181023 see CAS RNs attached to abstract & J. CHEM. SOC. C (1971), (3), 425-31 ,</p>	8
A	<p>GB 2 277 090 A (SHELL INT RESEARCH) 19 October 1994 (1994-10-19) abstract; claims</p>	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/06565

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9202513	A	20-02-1992	WO JP 9202513 A1 6501926 T	20-02-1992 03-03-1994
WO 8500603	A	14-02-1985	US 4504482 A AU 3154184 A DK 132285 A EP 0150205 A1 FI 851167 A ,B, IL 72411 A IT 1213193 B JP 60502003 T NO 851113 A NZ 208817 A PH 19913 A WO 8500603 A1 ZA 8405424 A	12-03-1985 04-03-1985 22-03-1985 07-08-1985 22-03-1985 29-04-1988 14-12-1989 21-11-1985 20-03-1985 12-02-1988 14-08-1986 14-02-1985 27-02-1985
WO 0073278	A	07-12-2000	AU 5070800 A WO 0073278 A2	18-12-2000 07-12-2000
WO 9838171	A	03-09-1998	US 5935966 A AU 6666798 A WO 9838171 A1	10-08-1999 18-09-1998 03-09-1998
WO 9709311	A	13-03-1997	AU 708616 B2 AU 6743296 A BR 9610385 A CA 2230931 A1 CN 1202152 A CZ 9800684 A3 WO 9709311 A1 EP 0863875 A1 HU 9900926 A2 JP 11500447 T NO 980954 A PL 325425 A1 TR 9800409 T1 US 6051712 A US 6150526 A ZA 9607424 A	05-08-1999 27-03-1997 06-07-1999 13-03-1997 16-12-1998 14-10-1998 13-03-1997 16-09-1998 28-09-1999 12-01-1999 28-04-1998 20-07-1998 21-05-1998 18-04-2000 21-11-2000 07-03-1997
WO 9535293	A	28-12-1995	ES 2079323 A1 AU 2675095 A CA 2169050 A1 EP 0714896 A1 WO 9535293 A1 JP 9502205 T	01-01-1996 15-01-1996 28-12-1995 05-06-1996 28-12-1995 04-03-1997
WO 9111441	A	08-08-1991	DE 4003012 A1 CA 2075198 A1 WO 9111441 A1 EP 0513069 A1 IE 910362 A1 JP 2995089 B2 JP 5504349 T NO 922978 A PT 96647 A US 5366657 A	08-08-1991 03-08-1991 08-08-1991 19-11-1992 28-08-1991 27-12-1999 08-07-1993 28-07-1992 31-10-1991 22-11-1994

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/06565

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 8903416	A	20-04-1989	DE 3733818 A1 WO 8903416 A1 EP 0334925 A1 JP 2501482 T	20-04-1989 20-04-1989 04-10-1989 24-05-1990
EP 0295370	A	21-12-1988	DE 3709618 A1 AT 92921 T DE 3883090 D1 EP 0295370 A1 JP 2787572 B2 JP 63258460 A KR 9612176 B1 NO 881282 A US 4906752 A	06-10-1988 15-08-1993 16-09-1993 21-12-1988 20-08-1998 25-10-1988 16-09-1996 26-09-1988 06-03-1990
EP 0284008	A	28-09-1988	DE 3709549 A1 AT 96460 T DE 3885128 D1 EP 0284008 A2 JP 2590191 B2 JP 63254183 A NO 881281 A US 4906401 A	06-10-1988 15-11-1993 02-12-1993 28-09-1988 12-03-1997 20-10-1988 26-09-1988 06-03-1990
EP 0257605	A	02-03-1988	AU 602913 B2 AU 7741287 A BR 8704368 A DK 443287 A EP 0257605 A1 TR 23028 A US 4847258 A JP 63068572 A ZA 8706303 A	01-11-1990 03-03-1988 19-04-1988 27-02-1988 02-03-1988 10-02-1989 11-07-1989 28-03-1988 25-05-1988
US 4438117	A	20-03-1984	NONE	
JP 4300871	A	23-10-1992	NONE	
DD 145913	Z		NONE	
GB 2277090	A	19-10-1994	NONE	